

---

---

**Optics and photonics — Preparation  
of drawings for optical elements and  
systems —**

**Part 12:  
Aspheric surfaces —  
AMENDMENT 1**

*Optique et photonique — Préparation des dessins pour éléments et  
systèmes optiques —*

*Partie 12: Surfaces asphériques —*

*AMENDEMENT 1*





**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2013

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 749 09 47  
E-mail [copyright@iso.org](mailto:copyright@iso.org)  
Web [www.iso.org](http://www.iso.org)

Published in Switzerland

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2. [www.iso.org/directives](http://www.iso.org/directives)

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received. [www.iso.org/patents](http://www.iso.org/patents)

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

The committee responsible for this document is ISO/TC 172, *Optics and photonics*, Subcommittee SC 1, *Fundamental standards*.

STANDARDSISO.COM : Click to view the full PDF of ISO 10110-12:2007/Amd 1:2013

# Optics and photonics — Preparation of drawings for optical elements and systems —

## Part 12: Aspheric surfaces

### AMENDMENT 1

Page iv, Foreword

In the list of part titles of ISO 10110, update the part title of Part 8 to read:

— *Surface texture; roughness and waviness*

Page 2, 3.1.2

Number the NOTE to become NOTE 1 and add the following NOTE 2 after NOTE 1:

NOTE 2 In this case, “left” and “right” presume  $z$  is increasing from left to right. When the  $Z$ -axis is reversed as a result of a reflection (a  $180^\circ$  rotation about the  $Y$ -axis), the sign convention for radius and sagitta is also reversed. This is discussed further in 3.3.2.3.

Page 5, 3.3.2.1

Immediately after Equation (12), i.e. before the sentence about Schmidt surface, insert the following new paragraph:

The first order and second order aspheric terms,  $A_1h$  and  $A_2h^2$ , may be added to Equation (12).

Page 6, 3.3.2.3

Add the following new subclause after 3.3.2.3:

#### 3.3.2.4 Combined surface based on an orthogonal polynomial

A surface of higher order can also be generated by combining a spherical surface [Equation (5)] with a set of orthogonal polynomials of the following type possessing orthonormal derivatives (see Reference [3]).

$$z = f(x, y) + f_1(x, y) \quad (17)$$

$$f_1(x, y) = \frac{\left(\frac{h}{h_0}\right)^2 \left[1 - \left(\frac{h}{h_0}\right)^2\right]}{\sqrt{1 - \left(\frac{h}{R}\right)^2}} \left[ A_0 Q_0\left(\frac{h^2}{h_0^2}\right) + A_1 Q_1\left(\frac{h^2}{h_0^2}\right) + \dots \right] \quad (18)$$

where  $h_0$  marks the upper limit of  $h$ . The description  $z$  is valid for  $0 \leq h \leq h_0$  only.  $R$  is the radius of curvature that intersects the surface at  $h_0$ .

For a detailed description of the recursion formulae, see Annex B.

NOTE Another description that combines a conic with a set of orthogonal polynomials, orthogonal in amplitude, is also permitted [see Reference [4], Equations (2) to (3) and (8) to (9)].

Page 14, Annex A

In the table, add a new row for "Sphere", under "Surfaces rotationally symmetric about Z-axis". For the sake of clarity, the amended table of Annex A is given hereafter in its entirety.

| Class                                                       | Basic surface                                    | Basic equation $f(x,y) =$                                                                                                                              | Power series $f_1(x,y) =$ [for toric surfaces, $g_1(x)$ ]                                                                                                                                                                                     |
|-------------------------------------------------------------|--------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Non-rotationally-symmetric surfaces                         | Ellipsoid<br>Hyperboloid<br>Paraboloid           | $\frac{x^2}{R_X} + \frac{y^2}{R_Y}$<br>$1 + \sqrt{1 - (1 + \kappa_X) \left( \frac{x}{R_X} \right)^2 - (1 + \kappa_Y) \left( \frac{y}{R_Y} \right)^2}$  | $A_4 x^4 + B_4 y^4 + A_6 x^6 + B_6 y^6 + \dots$<br>$\dots C_3  x ^3 + \dots + D_3  y ^3 + \dots$                                                                                                                                              |
| $R_X \neq R_Y^a$<br>$\kappa_X \neq \kappa_Y$                | Cone ( $a \neq b$ )                              | $c \sqrt{\frac{x^2}{a^2} + \frac{y^2}{b^2}}$                                                                                                           |                                                                                                                                                                                                                                               |
| $A_{2i} \neq B_{2i}$<br>$C_{2i-1} \neq D_{2i-1}$            | Cylinder                                         | $\frac{u^2}{R_U \left[ 1 + \sqrt{1 - (1 + \kappa_U) \left( \frac{u}{R_U} \right)^2} \right]}$                                                          | $A_4 x^4 + A_6 x^6 + \dots + C_3  x ^3$<br>for $u = x$<br>$B_4 y^4 + B_6 y^6 + \dots + D_3  y ^3$<br>for $u = y$                                                                                                                              |
| Surfaces rotationally symmetric about Z axis                | Ellipsoid<br>Hyperboloid<br>Paraboloid<br>Sphere | $\frac{h^2}{R \left[ 1 + \sqrt{1 - (1 + \kappa) \left( \frac{h}{R} \right)^2} \right]}$                                                                | $A_3 h^3 + A_4 h^4 + A_5 h^5 + \dots$                                                                                                                                                                                                         |
| $R_X = R_Y = R$<br>$\kappa_X = \kappa_Y = \kappa$           | Cone ( $a = b$ )                                 | $\frac{c}{a} h$                                                                                                                                        | where<br>$h = \sqrt{x^2 + y^2}$                                                                                                                                                                                                               |
| $h^2 = x^2 + y^2$                                           | Plane (Schmidt surface)<br>Sphere                | 0<br>$\frac{h^2}{R \left[ 1 + \sqrt{1 - \frac{h^2}{R^2}} \right]}$                                                                                     | $\frac{\left( \frac{h}{h_0} \right)^2 \left[ 1 - \left( \frac{h}{h_0} \right)^2 \right]}{\sqrt{1 - \left( \frac{h}{R} \right)^2}} \left[ A_0 Q_0 \left( \frac{h^2}{h_0^2} \right) + A_1 Q_1 \left( \frac{h^2}{h_0^2} \right) + \dots \right]$ |
| Surfaces of revolution; not coincident with coordinate axis | Toric surface                                    | $f(x,y) = R_Y \mp \sqrt{[R_Y - g(x)]^2 - y^2}$<br>$g(x) = \frac{x^2}{R_X \left[ 1 + \sqrt{1 - (1 + \kappa_X) \left( \frac{x}{R_X} \right)^2} \right]}$ | $g_1(x) = A_4 x^4 + A_6 x^6 + \dots + C_3  x ^3 + C_5  x ^5 + \dots$                                                                                                                                                                          |
| a If at least one of these inequalities is valid.           |                                                  |                                                                                                                                                        |                                                                                                                                                                                                                                               |

Page 15, new Annex B

Add the following annex:

**Annex B**  
(informative)

**Description of an orthogonal polynomial**

$$Q_{m+1}\left(\frac{h^2}{h_0^2}\right) = \frac{P_{m+1}\left(\frac{h^2}{h_0^2}\right) - g_m Q_m\left(\frac{h^2}{h_0^2}\right) - k_{m-1} Q_{m-1}\left(\frac{h^2}{h_0^2}\right)}{l_{m+1}} \quad (\text{B.1})$$

$$\text{starting with } Q_0\left(\frac{h^2}{h_0^2}\right) = 1 \text{ and } Q_1\left(\frac{h^2}{h_0^2}\right) = \frac{13 - 16\left(\frac{h^2}{h_0^2}\right)}{\sqrt{19}}$$

$$P_{m+1}\left(\frac{h^2}{h_0^2}\right) = \left(2 - 4\frac{h^2}{h_0^2}\right)P_m\left(\frac{h^2}{h_0^2}\right) - P_{m-1}\left(\frac{h^2}{h_0^2}\right) \quad (\text{B.2})$$

$$\text{starting with } P_0\left(\frac{h^2}{h_0^2}\right) = 2 \text{ and } P_1\left(\frac{h^2}{h_0^2}\right) = 6 - 8\left(\frac{h^2}{h_0^2}\right).$$

The following auxiliary polynomials (B.3), (B.4) and (B.5) have to be solved in the order given here and are valid for  $m \geq 2$ .

$$k_{m-2} = \frac{-m(m-1)}{2 l_{m-2}} \quad (\text{B.3})$$

$$g_{m-1} = \frac{-(1 + g_{m-2}k_{m-2})}{l_{m-1}} \quad (\text{B.4})$$

$$l_m = \left[ m(m+1) + 3 - g_{m-1}^2 - k_{m-2}^2 \right]^{1/2} \quad (\text{B.5})$$

$$\text{starting with } g_0 = -\frac{1}{2}, \quad l_0 = 2 \text{ and } l_1 = \frac{1}{2}\sqrt{19}.$$

Based on the recursion the first six  $Q$ s are the following:

$$Q_0\left(\frac{h^2}{h_0^2}\right) = 1$$

$$Q_1\left(\frac{h^2}{h_0^2}\right) = \frac{13 - 16\left(\frac{h^2}{h_0^2}\right)}{\sqrt{19}}$$

$$Q_2\left(\frac{h^2}{h_0^2}\right) = \sqrt{\frac{2}{95}} \left[ 29 - 4\left(\frac{h^2}{h_0^2}\right) \left( 25 - 19\left(\frac{h^2}{h_0^2}\right) \right) \right]$$

$$Q_3\left(\frac{h^2}{h_0^2}\right)=\sqrt{\frac{2}{2545}}\left\{207-4\left(\frac{h^2}{h_0^2}\right)\left[315-\left(\frac{h^2}{h_0^2}\right)\left(577-320\left(\frac{h^2}{h_0^2}\right)\right)\right]\right\}$$

$$Q_4\left(\frac{h^2}{h_0^2}\right)=\frac{1}{3\sqrt{131831}}\left\{7737-16\left(\frac{h^2}{h_0^2}\right)\left[4653-2\left(\frac{h^2}{h_0^2}\right)\left[7381-8\left(\frac{h^2}{h_0^2}\right)\left(1168-509\left(\frac{h^2}{h_0^2}\right)\right)\right]\right]\right\}$$

$$Q_5\left(\frac{h^2}{h_0^2}\right)=\frac{1}{3\sqrt{6632213}}\left\{66657-32\left(\frac{h^2}{h_0^2}\right)\left[28338-\left(\frac{h^2}{h_0^2}\right)\left[135325-8\left(\frac{h^2}{h_0^2}\right)\left[35884-\left(\frac{h^2}{h_0^2}\right)\left(34661-12432\left(\frac{h^2}{h_0^2}\right)\right)\right]\right]\right]\right\}$$

Page 15, Bibliography

Add the following bibliography references:

- [2] KROSS, J., OERTMANN, F.-W., SCHUHMAN, R., *On aspherics in optical system*, SPIE Proceedings 656 (1986)
- [3] FORBES, G.W., Robust, efficient computational methods for axially symmetric optical aspheres, Opt. Express 18, 19700-19712 (2010)
- [4] FORBES, G.W., Shape specification for axially symmetric optical surfaces, Opt. Express 15, 5218-5226 (2007)